ELSEVIER

Volume 514, Numbers 1–2, 17 May 1996

Journal of Organometallic Chemistry

Contents

Regular Papers

Infrared and nuclear magnetic resonance spectroscopic studies of the structure and dynamics of allylic magnesium compounds
E.A. Hill, W.A. Boyd, H. Desai, A. Darki and L. Bivens (Milwaukee, WI, USA) 1

Photoinduced rearrangements of ring-coupled bis(cyclopentadienyl)rutheniumdicarbonyl) compounds of the form $Ru_2(CO)_4(\eta^5,\eta^5\text{-}C_5H_4LC_5H_4)$ (L = CH_2–, $(CH_3)_2C$–, C_2H_4– and $(CH_3)_2Si$–). The molecular structure of $[Ru(CO)_2][Ru(CO)_2Cl](\eta^5,\eta^5{:}\eta^1\text{-}C_5H_4CH_2C_5H_3)$
T.E. Bitterwolf (Moscow, ID, USA), J.E. Shade, J.A. Hansen (Annapolis, MD, USA) and A.L. Rheingold (Newark, DE, USA). 13

Deposition of silicon carbide films by plasma enhanced chemical vapour deposition
J. Ramirez, H. Suhr (Tubingen, Germany), L. Szepes, L. Zanathy and A. Nagy (Budapest, Hungary) 23

Cyclosiloxane sandwich complexes of a lanthanide metal: $Na_6\{[(C_6H_3SiO_2)_3]_2Nd_4(\mu_4\text{-}O)\}$
O.I. Shchegolikhina, Yu.A. Pozdniakova, S.V. Lindeman, A.A. Zhdanov (Moscow, Russia), R. Psaro, R. Ugo (Milano, Italy), G. Gavioli, R. Battistuzzi, M. Borsari, T. Rüffer, C. Zucchi and G. Pályi (Modena, Italy) 29

Diphosphirane and 1,3-diphosphaallyl anion: geometry and bonding
M. Gouygou (Paris, France), M. Veith (Saarbrücken, Germany), C. Couret, J. Escudié (Toulouse, France), V. Huch (Saarbrücken, Germany) and M. Koenig (Toulouse, France) 37

Unverbrückte Indenylkomplexe mit Metallen der vierten Gruppe als Katalysatoren bei der homogenen Olefinpolymerisation
M.A. Schmid, H.G. Alt und W. Milius (Bayreuth, Deutschland) 45

Pentamethylcyclopentadienyl halo– and alkyl–alkoxo tantalum(V) complexes. Crystal structure of $TaCp^*(CH_2SiMe_3)_2\{\eta^2\text{-}O(2\text{-}CH_2\text{–}6\text{-}MeC_6H_3)\}$
I. De Castro, M.V. Galakhov, M. Gómez, P. Gómez-Sal, A. Martin and P. Royo (Alcalá de Henares, Spain) 51

Le diménitylnéopentylsilène, un silène stable et facile à préparer synthèse et aspects de sa réactivité
G. Delpon-Lacaze, C. De Battisti et C. Couret (Toulouse, France) 59

Reaction of dihydrosilanes with ethyl diazoacetate: synthesis of 3,3-disubstituted 3-silaglutarates
J.-P. Barnier and L. Blanco (Orsay, France) 67

Synthesis and X-ray crystal structures of alkaline-earth metallocenes with pendant substituents
M.L. Hays, T.P. Hanusa (Nashville, TN, USA) and T.A. Nile (Greensboro, NC, USA). 73

Metallorganische Verbindungen des Kupfers XI. Synthese und Struktur von zwei vierkernigen Alkin-μ^2-Halogeno-μ^4-Oxo–Kupfer(I)-Komplexen; Alkin = 3,3,6,6-Tetramethyl-1-thia-4-cycloheptin
G. Gröger, F. Olbrich, E. Weiss und U. Behrens (Hamburg, Deutschland) 81

Crystal structure of μ_3-hydridododecacarbonyltriosmium dioxooctafluoro-μ_3-fluoroditungsten
M.C. Crossman, J. Fawcett, E.G. Hope and D.R. Russell (Leicester, UK) 87

Synthesis and characterization of tert-butylcyclopentadienyl derivatives of hafnium
T. Cuenca, C. Montejano and P. Royo (Alcalá de Henares, Spain). 93

Efficient catalytic reduction of ketones with formic acid and ruthenium complexes
N. Menashe, E. Salant and Y. Shvo (Tel Aviv, Israel) 97

$[CpTi(\mu_3\text{-}S)_3\{Rh(tfbb)\}_3]$: An early–late heterometallic complex as molecular model for the deactivation of metal sulphide catalysts
R. Atencio, M.A. Casado, M.A. Ciriano, F.J. Lahoz, J.J. Pérez-Torrente (Zaragoza, Spain), A. Tiripicchio (Parma, Italy) and L.A. Oro (Zaragoza, Spain). 103

Structure and properties of dimer, trimer and tetramer aggregates of methyltrioxorhenium (MTO): an ab initio study
S. Köstlmeier, G. Pacchioni (Milan, Italy), W.A. Herrmann and N. Rösch (Milan, Garching, Italy, Germany) 111

Dinuclear rhodium–rhodium and rhodium–palladium complexes with different bridging ligands. Crystal structure of $[Rh_2(\mu\text{-}SPPh_2)(\mu\text{-}Cl)(COD)_2]$
M.T. Pinillos, M.P. Jarauta, A. Elduque, F.J. Lahoz and L.A. Oro (Zaragoza, Spain) 119

Coordination chemistry of 1,1′-bis(2-pyridyl)ferrocene in palladium(II) and platinum(II) complexes and reaction of (BPF)Pd(Me)Cl with CO. X-ray structure of (1,1′-bis(2-pyridyl)ferrocene)Pd(Me)Cl and (1,1′-bis(2-pyridyl)ferrocene)$[PtCl_2(CH_2=CH_2)]_2$
J.G.P. Delis, P.W.N.M. Van Leeuwen, K. Vrieze (Amsterdam, Netherlands), N. Veldman, A.L. Spek, J. Fraanje and K. Goubitz (Utrecht, Amsterdam, Netherlands) 125

Synthesis, redox chemistry and solid state structure of di- and trinuclear ferrocenyl substituted *N*-alkylpyrazolyl pyridine complexes
W.R. Thiel, T. Priermeier, D.A. Fiedler (Garching, Germany), A.M. Bond (Clayton, Australia) and M.R. Mattner (Garching, Germany) 137

Synthesis of $(p\text{-TolSb})_n$ by reaction of Cp_2Co with p-$TolSbCl_2$, crystal structure of $[Cp_2Co][p\text{-}TolSbCl_3]$, and electroreduction of p-$TolSbCl_2$
H.J. Breunig, K.H. Ebert, J. Probst (Bremen, Germany), Y. Mourad and Y. Mugnier (Dijon, France) 149

A novel heterobimetallic rotation system: characterization of palladium(II) complexes with 2-pyridylethylferrocenecarboxamides
T. Moriuchi, I. Ikeda and T. Hirao (Osaka, Japan) 153

Stereospecific syntheses of functionalised iron alkyl complexes with a stereogenic metal center from an iron formyl precursor
M.A. Guillevic, P. Brégaint and C. Lapinte (Rennes, France) 157

The crystal structure of di-2-aminopyrimidinium tetraphenyldichloro(μ-oxo)distannate
N. Kourkoumelis (Ioannina, Greece), A. Hatzidimitriou (Toulouse, France) and D. Kovala-Demertzi (Ioannina, Greece). 163

Different coordination modes of the 1,1,1-tris(diphenylphosphinomethyl)ethane ligand in gold(I) and gold(III) complexes
E.J. Fernández (Logroño, Spain), M.C. Gimeno, A. Laguna (Zaragoza, Spain), M. Laguna, J.M. López-de-Luzuriaga and E. Olmos (Logroño, Spain) . . . 169
Ionic complexes $[Au_2(P-P)_2][Co(CO)_4]_2$ as intermediates in the synthesis of compounds containing a Co_2Au_2 core
A. Pons, O. Rossell, M. Seco, X. Solans and M. Font-Bardía (Barcelona, Spain) . . . 177
Synthesis and molecular structures of di- and trinuclear molybdenum complexes containing pyridine-2-thiolato (pyS) ligand
Y. Shi, S. Lu, H. Guo (Dalian, People's Republic of China), Q. Wu (Fuzhou, People's Republic of China) and N. Hu (Changchun, People's Republic of China) . . . 183
The solid state structures of *ortho*-lithio-β-(*N*,*N*-dimethylamino)ethoxybenzene and the corresponding sodium compound: a new type of organolithium tetramer and a first example of an organosodium hexamer
R. Den Besten, M.T. Lakin, N. Veldman, A.L. Spek and L. Brandsma (Utrecht, Netherlands) . . . 191
Tri- and binuclear ruthenium carbonyl complexes containing bridging ligands derived from *ortho*-substituted anilines. A comparative study of their syntheses using $RuCl_3 \cdot nH_2O$ or $[Ru_3(CO)_{12}]$ as starting materials
J.A. Cabeza, M.A. Martínez-García, V. Riera, S. García-Granda and J.F. Van der Maelen (Oviedo, Spain) . . . 197
Synthesis and characterization of triphenyl- and tri-n-butyltin pentafluorobenzoates, -phenylacetates and -cinnamates. X-ray structure determination of tri-n-butyltin pentafluorocinnamate
R. Willem, A. Bouhdid, M. Biesemans, J.C. Martins (Brussels, Belgium), D. De Vos (Haarlem, Netherlands), E.R.T. Tiekink (Adelaide, Australia) and M. Gielen (Brussels, Belgium) . . . 203
Molecular structure of 2,2′-thiobis(4-methyl-6-tert-butylphenoxy) titanium diisopropoxy. Influence of titanium–sulfur interaction on coordination geometry
L. Porri, A. Ripa, P. Colombo, E. Miano, S. Capelli and S.V. Meille (Milan, Italy) . . . 213
Synthesis and structure of titanium(III) tweezer complexes with embedded alkali metal ions: $[(\eta^5\text{-}C_5HMe_4)_2Ti(\eta^1\text{-}C{\equiv}SiMe_3)_2]^-M^+$ (M = Li, Na, K or Cs)
V. Varga, J. Hiller, M. Polášek, U. Thewalt and K. Mach (Prague, Czech Republic) . . . 219
Regio- and stereoselective a^4-umpolung reactions of α,β-unsaturated esters to 1,6-dicarbonyl compounds by addition of enantiopure nucleophiles to racemic tetracarbonyl (η^3-allyl)iron(1 +) complexes
D. Enders (Aachen, Germany), P. Fey (Wuppertal, Germany), T. Schmitz (Berlin, Germany), B.B. Lohray (Hyderabad, India) and B. Jandeleit (Aachen, Germany) . . . 227
The structural behaviour of 2,2-functionally disubstituted triorganotin chlorides
F. Richter (Leverkusen, Germany), M. Dargatz (Geelong, Australia), H. Hartung, D. Schollmeyer and H. Weichmann (Halle, Germany) . . . 233
Synthesis of mono-phosphido, mono-dppm bridged Pd(I) cationic dimers. Structure and fluxionality of $[Pd_2(\mu\text{-}P^tBu_2)(\mu\text{-dppm})(\eta^1\text{-dppm})_2]CF_3SO_3$ (dppm = bis-diphenylphosphinomethane)
P. Leoni, M. Pasquali, G. Pieri (Pisa, Italy) and U. Englert (Aachen, Germany) . . . 243
Syndiospezifische Polymerisation von Propylen: 3-, 4-, 3,4- und 4,5-substituierte Zirkonocenkomplexe des Typs $(C_{13}H_{8-n}R_nCR'_2C_5H_4)ZrCl_2$ (n = 1, 2; R = Alkyl, Aryl; R′ = Me, Ph)
H.G. Alt, R. Zenk und W. Milius (Bayreuth, Deutschland) . . . 257
Copolymerization of ethylene and carbon monoxide by phosphinite-modified palladium catalysts
W. Keim and H. Maas (Aachen, Germany) . . . 271
The mechanism of the isomerization of organosilicon cyanates to isocyanates
A.I. Almansour (Riyadh, Saudi Arabia), G.A. Ayoko and C. Eaborn (Brighton, UK) . . . 277
Indol-1-yl- und Pyrrol-1-yl-substituierte Verbindungen des Siliciums und Phosphors
A. Frenzel, M. Gluth, R. Herbst-Irmer und U. Klingebiel (Göttingen, Deutschland) . . . 281
Synthesis and characterization of novel neutral σ-alkynylcarbonylnitrosyl molybdenum(0) and tungsten(0) complexes. Crystal structure of $[Mo(C{\equiv}C^nPr)(CO)_2(dppe)(NO)]$ [dppe = 1,2-bis(diphenylphosphino)ethane]
M.P. Gamasa, J. Gimeno, L. Zhang (Oviedo, Spain), M. Lanfranchi and A. Tiripicchio (Parma, Italy) . . . 287

Author Index . . . 299

Subject Index . . . 301

The publisher encourages the submission of articles in electronic form thus saving time and avoiding rekeying errors. A leaflet describing our requirements is available from the publisher upon request.